

3150 Great Eastern Hwy Glen Forrest

Western Australia 6071

p: +61 8 9298 8111 F: +61 8 9298 8700

Email: info@specialtyfeeds.com

Diet **SF00-219**

21% Fat, 0.15% Cholesterol Semi-Pure Rodent Diet

A semi-pure high fat diet formulation for laboratory rats and mice based formulated to mimic a "Western fast food diet". This diet was originally formulated to be similar to *Harlan Teklad* TD88137 or *Research Diets* Western Diet D12079B. Some modifications have been made to the original formulation to suit locally available raw materials.

 This Diet was developed to generate arteriosclerotic lesions in a range of susceptible mice strains.

| Calculated Nutritional Parameters | | |
|--|-----------------------|--|
| Protein | 19.0% | |
| Total Fat | 21.0% | |
| Total digestible carbohydrate as defined by FSANZ Standard 1.2.8 | 47.3% | |
| Crude Fibre | 4.70% | |
| AD Fibre | 4.70% | |
| Net Metabolisable Energy | 17.8 MJ / Kg | |
| Digestible Energy | 19 MJ / Kg | |
| % Total calculated energy from lipids | 41.0% DE 43.0% NME | |
| % Total calculated energy from protein | 17.0% DE 14.0% NME | |

Diet Form and Features

- Semi pure high fat diet. 12 mm diameter pellets.
- Pack size 2 Kg compostable cardboard trays, vacuum packed in oxygen impermeable plastic bags, under nitrogen.
 Bags are packed into cardboard cartons to protect them during transit. Smaller pack quantity on request.
- Diet suitable for irradiation but not suitable for autoclave.
- Lead time 2 weeks for non-irradiation or 4 weeks for irradiation.

| Ingredients | |
|--------------------------------|-----------|
| Casein (Acid) | 195 g/Kg |
| Sucrose | 341 g/Kg |
| Clarified Butter (Ghee) | 210 g/Kg |
| Cellulose | 50 g/Kg |
| Wheat Starch | 154 g/Kg |
| L Methionine | 3.0 g/Kg |
| Calcium Carbonate | 17.1 g/Kg |
| Sodium Chloride | 2.6 g/Kg |
| AIN93 Trace Minerals | 1.4 g/Kg |
| Potassium Citrate | 2.6 g/Kg |
| Potassium Dihydrogen Phosphate | 6.9 g/Kg |
| Potassium Sulphate | 1.6 g/Kg |
| Choline Chloride (75%) | 2.5 g/Kg |
| SF00-219 Vitamins | 10 g/Kg |
| Cholesterol | 1.5 g/Kg |
| Oxicap E2 | 0.04 g/Kg |

Ppenness Agreement
Animal Research and Teaching in Australia
UPPORTER

RUSTED TRADER



| Calculated Amino Acids as I | alculated Amino Acids as Fed | |
|-----------------------------|------------------------------|--|
| Valine | 1.00% | |
| Leucine | 1.60% | |
| Isoleucine | 0.87% | |
| Threonine | 0.59% | |
| Methionine | 0.79% | |
| Cysteine | 0.11% | |
| Lysine | 1.50% | |
| Phenylalanine | 0.89% | |
| Tyrosine | 1.10% | |
| Tryptophan | 0.17% | |
| Arginine | 0.49% | |
| Histidine | 0.20% | |

| | Calculated Total Vitamins as Fed | |
|-----|----------------------------------|--------------|
| | Vitamin A (Retinol) | 11 650 IU/Kg |
| | Vitamin D (Cholecalciferol) | 1 100 IU/Kg |
| | Vitamin E (a Tocopherol acetate) | 64 mg/Kg |
| | Vitamin K (Menadione) | 12.5 mg/Kg |
| | Vitamin C (Ascorbic acid) | 700 mg/Kg |
| | Vitamin B1 (Thiamine) | 11 mg/Kg |
| | Vitamin B2 (Riboflavin) | 11 mg/Kg |
| | Niacin (Nicotinic acid) | 50 mg/Kg |
| | Vitamin B6 (Pryridoxine) | 11 mg/Kg |
| | Pantothenic Acid | 34 mg/Kg |
| | Biotin | 200 ug/Kg |
| _ | Folic Acid | 1 mg/Kg |
| | Inositol | 55 mg/Kg |
| | Vitamin B12 (Cyancobalamin) | 18 ug/Kg |
| | Choline | 4 000 mg/Kg |
| | | |
| - 1 | | |

| Calculated Total Minerals as Fed | |
|----------------------------------|------------|
| Calcium | 0.61% |
| Phosphorous | 0.32% |
| Magnesium | 0.07% |
| Sodium | 0.12% |
| Chloride | 0.16% |
| Potassium | 0.39% |
| Sulphur | 0.22% |
| Iron | 73 mg/Kg |
| Copper | 7.0 mg/Kg |
| lodine | 0.2 mg/Kg |
| Manganese | 19 mg/Kg |
| Cobalt | No data |
| Zinc | 45 mg/Kg |
| Molybdenum | 0.15 mg/Kg |
| Selenium | 0.3 mg/Kg |
| Cadmium | No data |
| Chromium | 1.0 mg/Kg |
| Fluoride | 1.0 mg/Kg |
| Lithium | 0.1 mg/Kg |
| Boron | 2.0 mg/Kg |
| Nickel | 0.5 mg/Kg |
| Vanadium | 0.1 mg/Kg |

| _ | | |
|---|--|---------|
| | Calculated Fatty Acid Composition as Fed | |
| | Saturated Fats C12:0 or less | 1.70% |
| | Myristic Acid 14:0 | 2.50% |
| | Palmitic Acid 16:0 | 6.90% |
| | Stearic Acid 18:0 | 2.30% |
| | Palmitoleic Acid 16:1 | 0.21% |
| | Oleic Acid 18:1 | 5.50% |
| | Gadoleic Acid 20:1 | No data |
| | Linoleic Acid 18:2 n6 | 0.21% |
| | a Linolenic Acid 18:3 n3 | 0.17% |
| | Arachadonic Acid 20:4 n6 | Trace |
| | Total n3 | 0.35% |
| | Total n6 | 0.25% |
| | Cholesterol | 0.15% |
| | Total Mono Unsaturated Fats | 5.90% |
| | Total Polyunsaturated Fats | 0.80% |
| | Total Saturated Fats | 13.60% |
| | Denness Agreement A Animal Research and Teaching in Australia | |

Calculated data uses information from typical raw material composition. **Diet post treatment by irradiation or autoclave could change these parameters**. It could be expected that individual batches of diet will vary from this figure. We are happy to provide full calculated nutritional information for all of our products, however we would like to emphasise that these diets have been specifically designed for manufacture by Specialty Feeds.

SUPPORTER